


AMENDMENTS TO THE CLAIMS

Listing of Claims


The following listing of claims will replace all prior versions and listing of claims in the application.

1. (currently amended) A compound of Formula I



or a pharmaceutically acceptable [[salts]] salt thereof wherein:

X is



Y is selected from the group consisting of N-R¹, O, and S;

A is N or C;

~~R¹ is selected from the group consisting of H, alkyl, aryl, hydroxy, alkoxy, cyano, nitro, amino, alkenyl, alkynyl, amido, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxy carbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxy carbonyl, alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxyl, haloalkyl, cyano, nitro, carboxyl, amino, alkoxy, aryl or aryl optionally substituted with one or more~~

~~halogen, haloalkyl, lower alkyl, alkoxy, cyano, alkylsulfonyl, alkylthio, nitro, carboxyl, amino, hydroxyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, hydroxy, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, cyano, nitro, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, carboxyl derivatives, amino, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycle, and monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, sulfonic acid, sulfonamide, aryl or fused aryl; or~~


R¹ taken together with R⁸ forms a 4-12 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, alkoxy, halo, phenyl, amino, carboxyl or carboxyl ester, and fused phenyl; or

R¹ taken together with R⁸ forms a 5 membered heteroaromatic ring optionally substituted with one or more substituent selected from lower alkyl, phenyl and hydroxy; or

R¹ taken together with R⁸ forms a 5 membered heteroaromatic ring fused with a phenyl group;


~~R⁸ (when not taken together with R¹) and R⁹ are independently~~ is selected from the group consisting of H, alkyl, alkenyl, alkynyl, aralkyl, amino, alkylamino, hydroxy, alkoxy, arylamino, amido, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylthiocarbonyl, acyloxymethoxycarbonyl, cycloalkyl, bicycloalkyl, aryl, acyl, benzoyl, alkyl optionally substituted with one or more substituent selected from lower alkyl, halogen, hydroxy, haloalkyl, cyano, nitro, carboxyl derivatives, amino, alkoxy, thio, alkylthio, sulfonyl, aryl, aralkyl, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido,

acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethyl, sulfonyl, alkylsulfonyl, haloalkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, fused monocyclic heterocycles, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, carboxyl derivatives, aryloxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles, or fused monocyclic heterocycles, monocyclic heterocycles optionally substituted with one or more substituent selected from halogen, haloalkyl, lower alkyl, alkoxy, aryloxy, amino, nitro, hydroxy, carboxyl derivatives, cyano, alkylthio, alkylsulfonyl, aryl, fused aryl, monocyclic and bicyclic heterocyclicalkyls, -SO₂R¹⁰ wherein R¹⁰ is selected from the group consisting of alkyl, aryl and monocyclic heterocycles, all optionally substituted with one or more substituent selected from the group consisting of halogen, haloalkyl, alkyl, alkoxy, cyano, nitro, amino, acylamino, trifluoroalkyl, amido, alkylaminosulfonyl, alkylsulfonyl, alkylsulfonylamino, alkylamino, dialkylamino, trifluoromethylthio, trifluoroalkoxy, trifluoromethylsulfonyl, aryl, aryloxy, thio, alkylthio, and monocyclic heterocycles; and



NR⁸ and R⁹ taken together form a 4-12 membered mononitrogen containing monocyclic or bicyclic ring optionally substituted with one or more substituent selected from lower alkyl, carboxyl derivatives, aryl or hydroxy and wherein said ring optionally contains a heteroatom selected from the group consisting of O, N, and S;


or




wherein Y^1 is selected from the group consisting of alkyl, cycloalkyl, bicycloalkyl, aryl, monocyclic heterocycles, alkyl optionally substituted with aryl which can also be optionally substituted with one or more substituent selected from halogen, haloalkyl, alkyl, nitro, hydroxy, alkoxy, aryloxy, aryl, or fused aryl, aryl optionally substituted with one or more substituent selected from halogen, haloalkyl, hydroxy, alkoxy, aryloxy, aryl, fused aryl, nitro, methylenedioxy, ethylenedioxy, or alkyl, alkynyl, alkenyl, $-S-R^{11}$ and $-OR^{11}$ wherein R^{11} is selected from the group consisting of H, alkyl, aralkyl, aryl, alkenyl, and alkynyl, or R^{11} taken together with R^8 forms a 4-12 membered mononitrogen and monosulfur or monooxygen containing heterocyclic ring optionally substituted with lower alkyl, hydroxy, keto, phenyl, carboxyl, or carboxyl ester, and fused phenyl, or R^{11} taken together with R^8 is thiazole, oxazole, benzoxazole, or benzothiazole;

~~R^8 is defined as above;~~ or

Y^1 (when Y^1 is carbon) taken together with R^8 forms a 4-12 membered mononitrogen or dinitrogen containing ring optionally substituted with alkyl, aryl, keto or hydroxy; or



wherein R^1 and R^8 taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, phenyl, or carboxyl derivatives; and R^9 is selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxy carbonyl, alkylthiocarbonyl, arylthiocarbonyl, or acyloxymethoxy carbonyl; or



X is

wherein R¹ and R⁸ taken together form a 5-8 membered dinitrogen containing heterocycle optionally substituted with hydroxy, keto, phenyl, or alkyl; and


R⁹ are both selected from the group consisting of alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aryloxy carbonyl, haloalkylcarbonyl, haloalkoxy carbonyl, alkylthiocarbonyl, arylthiocarbonyl, and acyloxymethoxycarbonyl;

R², R³ and R⁴ are independently selected from one or more substituent selected from the group consisting of H, alkyl, hydroxy, alkoxy, aryloxy, halogen, haloalkyl, haloalkoxy, nitro, amino, alkylamino, acylamino, dialkylamino, cyano, alkylthio, alkylsulfonyl, carboxyl derivatives, trihaloacetamide, acetamide, aryl, fused aryl, cycloalkyl, thio, monocyclic heterocycles, fused monocyclic heterocycles, and X, wherein X is defined as above;

R⁵, R⁶, and R⁷ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, carboxyl derivatives, haloalkyl, cycloalkyl, monocyclic heterocycles, monocyclic heterocycles optionally substituted with alkyl, halogen, haloalkyl, cyano, hydroxy, aryl, fused aryl, nitro, alkoxy, aryloxy, alkylsulfonyl, arylsulfonyl, sulfonamide, thio, alkylthio, carboxyl derivatives, amino, amido, alkyl optionally substituted with one or more of halogen, haloalkyl, hydroxy, alkoxy, aryloxy, thio, alkylthio, alkynyl, alkenyl, alkyl, arylthio, alkylsulfoxide, alkylsulfonyl, arylsulfoxide, arylsulfonyl, cyano, nitro, amino, alkylamino, dialkylamino, alkylsulfonamide, arylsulfonamide, acylamide, carboxyl derivatives, sulfonamide, sulfonic acid, phosphonic acid derivatives, phosphinic acid derivatives, aryl, arylthio,

arylsulfoxide, or arylsulfone all optionally substituted on the aryl ring with halogen, alkyl, haloalkyl, cyano, nitro, hydroxy, carboxyl derivatives, alkoxy, aryloxy, amino, alkylamino, dialkylamino, amido, aryl, fused aryl, monocyclic heterocycles, and fused monocyclic heterocycles, monocyclic heterocyclic thio, monocyclic heterocyclic sulfoxide, and monocyclic heterocyclic sulfone, which can be optionally substituted with halogen, haloalkyl, nitro, hydroxy, alkoxy, fused aryl, or alkyl, alkylcarbonyl, haloalkylcarbonyl, and arylcarbonyl, aryl optionally substituted in one or more positions with halogen, haloalkyl, alkyl, alkoxy, aryloxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkylthio, thio, hydroxy, cyano, nitro, acyloxy, carboxyl derivatives, carboxyalkoxy, amido, acylamino, amino, alkylamino, dialkylamino, trifluoroalkoxy, trifluoromethylsulfonyl, alkylsulfonyl, sulfonic acid, sulfonamide, aryl, fused aryl, monocyclic heterocycles and fused monocyclic heterocycles; and all isomers, enantiomers, tautomers, racemates and polymorphs thereof.

2. (original) A compound according to claim 1




wherein:

R^5 and R^6 = H;

R^7 = H; alkyl, haloalkyl, carboxyalkyl, alkenyl, alkynyl, and phenyl, optionally substituted with one or more halogen atom.

3. (currently amended) A compound according to claim 1




I

wherein:

R^2 , R^3 , and R^4 are H, OH, or haloalkyl;






X is













Y is $N-R^1$ wherein ~~R^1 is selected from the group consisting of H, alkyl, aryl, hydroxy, alkoxy, cyano, and nitro; or~~





R^1 taken together with R^8 forms a 4-12 membered dinitrogen containing heterocycle optionally substituted with one or more substituent selected from the group consisting of lower alkyl, hydroxy, keto, alkoxy, halogen, phenyl, amino, carboxyl or carboxyl ester, and fused phenyl.


4. (currently amended) A compound selected from the group consisting of:












, and



and all the isomers, enantiomers, tautomers, racemates and polymorphs thereof.

5. (currently amended) A ~~pharmaceutical~~ composition useful in treating or preventing conditions mediated by the $\alpha_v\beta_3$ or $\alpha_v\beta_5$ integrin receptors in a mammal, the composition comprising a compound of Claim 1, ~~2, 3, or 4~~.

6. (withdrawn) A method of inhibiting a condition mediated by the $\alpha_v\beta_3$ or $\alpha_v\beta_5$ integrin comprising administering a therapeutically effective amount of a compound of Claim 1, 2, 3, or 4.

7. (withdrawn) The method according to Claim 6 wherein the condition treated is selected from the group consisting of tumor metastasis, solid tumor growth, angiogenesis, osteoporosis, humoral hypercalcemia of malignancy, smooth muscle cell migration, restenosis, atherosclerosis, macular degeneration, retinopathy, and arthritis.